



Contents lists available at [ScienceDirect](#)

Quaternary Science Reviews

journal homepage: www.elsevier.com/locate/quascirev



Corrigendum

Corrigendum to “A 550 ka record of aeolian activity near North West Cape, Australia: inferences from grain-size distributions and bulk chemistry of SE Indian Ocean deep-sea sediments” [Quat. Sci. Rev. 83 (2014) 83–94]



Jan-Berend W. Stuut ^{a, b, *}, Felix Temmesfeld ^a, Patrick De Deckker ^c

^a NIOZ – Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB Den Burg, The Netherlands

^b MARUM – Center for Marine Environmental Sciences, Bremen University, 28359 Bremen, Germany

^c Research School of Earth Sciences, The Australian National University, Canberra ACT 0200, Australia

The authors regret that when this paper was first published, the log-ratio of the elements Zr and Fe in Fig. 7C contained wrong values on the y-axis. The correct element ratio was already presented in Fig. 3C and is also presented below in the corrected version of Fig. 7. The correct data are available through the PANGAEA database at: <http://www.pangaea.de>.

DOI of original article: <http://dx.doi.org/10.1016/j.quascirev.2013.11.003>.

* Corresponding author. NIOZ – Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB Den Burg, The Netherlands.
E-mail address: jbstuut@nioz.nl (J.-B.W. Stuut).

<http://dx.doi.org/10.1016/j.quascirev.2014.07.012>

0277-3791/© 2014 Elsevier Ltd. All rights reserved.




Fig. 7. Downcore records of MD00-2361 which are interpreted in terms of palaeoenvironmental conditions in North West Cape. (A) Log Ti/Ca, a proxy for land-derived sedimentation; (B) Continental humidity, expressed as a log-ratio of the fluvial and aeolian end members; (C) Log(Zr/Fe), a proxy for continental humidity; (D) Total dust, expressed as the sum of the two aeolian end members. Marine Isotope Stages (M.I.S.) are indicated at the top.